



Mechanistic study of the ruthenium-catalyzed synthesis of amides from alcohols and amines

Makarov, Ilya; Fristrup, Peter; Madsen, Robert

Publication date:
2011

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):
Makarov, I., Fristrup, P., & Madsen, R. (2011). *Mechanistic study of the ruthenium-catalyzed synthesis of amides from alcohols and amines*. Abstract from EuCOMC XIX, Toulouse, France.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Mechanistic study of the ruthenium-catalyzed synthesis of amides from alcohols and amines


Ilya Makarov*, Peter Fristrup, Robert Madsen

Department of Chemistry, Technical University of Denmark, 2800 Lyngby, DENMARK

E-mail: ilym@kemi.dtu.dk

Webpage: <http://www.organic.kemi.dtu.dk/>

The amide bond is one of the most important linkages in organic chemistry and constitutes the key functional group in peptides, polymers and many pharmaceuticals [1]. Amides are often synthesized by coupling of carboxylic acids or its derivatives with amines. In spite of their efficiency these methods are not environmentally friendly since toxic reagents are used and a lot of waste is produced. Development of new synthetic methods which are both efficient and “green” continues to receive significant attraction. Recently in Madsen group a new method for amide synthesis has been discovered where alcohols and amines are coupled directly with the liberation of hydrogen [2, 3] (Scheme 1).



Scheme 1. Amide synthesis.

In the present work the mechanistic studies of this reaction have been carried out. The studies are divided into an experimental part and a theoretical part. In the experimental part reaction kinetics was studied. Para-substituted benzyl alcohols were used as substrates to elucidate the electronic influence of the para substituent on the reactivity, *i.e.* a Hammett study. By using DFT calculations with the B3LYP or M06 functionals Gibbs free energies and structures were determined for some metal complexes and transition states in the proposed catalytic cycle. From the experimental and theoretical studies the rate determining step will be established which is important to further optimize the reaction.

References:

- [1] J. W. Bode, *Curr. Opin. Drug Discov. Dev.* **2006**, 9, 765-775
- [2] L. U. Nordstrøm, H. Vogt, R. Madsen, *J. Am. Chem. Soc.* **2008**, 130, 17672-17673.
- [3] J. H. Dam, G. Osztrovszky, L. U. Nordstrøm, R. Madsen, *Chem. Eur. J.* **2010**, 16, 6820-6827.